

New oscillation criteria for linear matrix Hamiltonian systems

Yuan Gong Sun

Department of Mathematics, Qufu Normal University, Qufu, Shandong 273165, China

Received 30 November 2001

Submitted by R. O'Malley

Abstract

Some new oscillation criteria are established for the matrix linear Hamiltonian system $X' = A(t)X + B(t)Y$, $Y' = C(t)X - A^*(t)Y$ under the hypothesis: $A(t)$, $B(t) = B^*(t) > 0$, and $C(t) = C^*(t)$ are $n \times n$ real continuous matrix functions on the interval $[t_0, \infty)$, $(-\infty < t_0)$. These results are sharper than some previous results even for self-adjoint second order matrix differential systems. © 2003 Elsevier Science (USA). All rights reserved.

Keywords: Oscillation; Linear; Hamiltonian systems

1. Introduction

Consider the matrix linear Hamiltonian system

$$\begin{cases} X' = A(t)X + B(t)Y, \\ Y' = C(t)X - A^*(t)Y, \end{cases} \quad t \geq t_0, \quad (1.1)$$

where $A(t)$, $B(t)$, $C(t)$ are $(n \times n)$ -matrices, and B , C are Hermitian, i.e., $B^*(t) = B(t)$, $C^*(t) = C(t)$. By M^* we mean the conjugate transpose of the matrix M .

A Hermitian matrix $M \in C^{n \times n}$ is positive semidefinite (positive definite) if for all $u \in C^n$, $u \neq 0$, $u^*Mu \geq 0$ (> 0). A positive semidefinite (positive definite) Hermitian matrix M will be denoted by $M \geq 0$ ($M > 0$), and the usual ordering of the eigenvalues of M by $\lambda_1[M] \geq \lambda_2[M] \geq \dots \geq \lambda_n[M]$.

For any two solutions X_1, Y_1 and X_2, Y_2 of (1.1), the Wronskian $X_1^*(t)Y_2(t) - Y_1^*(t)X_2(t)$ is a constant matrix. In particular, for any solution X, Y of (1.1), $X^*(t)Y(t) -$

E-mail address: yugsun@263.net.

$Y^*(t)X(t)$ is a constant matrix. The solution X, Y of (1.1) is said to be conjoined if $X^*(t)Y(t) - Y^*(t)X(t) = 0$. A conjoined solution X, Y of (1.1) is said to be a conjoined basis of (1.1) if the rank of the $(2n \times n)$ -matrix $(X(t), Y(t))$ is n . A conjoined basis X and Y of (1.1) is said to be oscillatory on $[t_0, \infty)$ if $\det X(t)$ has arbitrarily large zeros. System (1.1) is said to be oscillatory on $[t_0, \infty)$ if one conjoined basis of (1.1) is oscillatory.

In the case when $A(t) \equiv 0$, $B(t) > 0$, (1.1) reduces to the second-order self-adjoint matrix differential system

$$(P(t)X')' + Q(t)X = 0 \quad (1.2)$$

with $P(t) = B^{-1}(t)$, $Q(t) = -C(t)$. The oscillation and nonoscillation of (1.1) or (1.2) have been extensively studied by many authors [1–13]. A discrete version of (1.2) is studied in [14]. However, all these results are given in the form of $\lim_{t \rightarrow \infty} \sup \lambda_1[\cdot] = +\infty$. In this paper we establish some new oscillation criteria which are presented in the form of $\lim_{t \rightarrow \infty} \sup \lambda_1[\cdot] > \text{const}$ for the system (1.1) by using one particular function $\Phi(t, s, r)$ defined as

$$\Phi(t, s, r) = (t-s)^\alpha (s-r)^\beta, \quad \text{where } \alpha, \beta > \frac{1}{2} \text{ are constants and } r \geq t_0. \quad (1.3)$$

Our results improve most known oscillation results even for the self-adjoint differential system (1.2). This can be seen by the examples given at the end of this paper.

2. Main results

Let $\phi(t)$ and $\theta(t)$ be positive, smooth and real-valued functions on $[t_0, \infty)$. Suppose that $B(t) > 0$. Let us make a change of unknown variables

$$U = \phi X, \quad V = \theta Y + \alpha B^{-1} X,$$

where

$$\alpha = \frac{\theta}{2} \left(\frac{\phi'}{\phi} - \frac{\theta'}{\theta} \right).$$

Then U and V satisfy a differential system

$$\begin{cases} U' = A(t)U + B_1(t)V + \frac{1}{2} \left(\frac{\phi'}{\phi} + \frac{\theta'}{\theta} \right) U, \\ V' = C_1(t)U - A^*(t)V + \frac{1}{2} \left(\frac{\phi'}{\phi} + \frac{\theta'}{\theta} \right) V, \end{cases} \quad (2.1)$$

where

$$B_1(t) = \frac{\phi(t)}{\theta(t)} B(t), \quad (2.2)$$

$$C_1(t) = \frac{\theta}{\phi} \left\{ C(t) + \frac{\alpha}{\theta} (B^{-1}(t)A(t) + A^*(t)B^{-1}(t)) + \left(\frac{\alpha}{\theta} B^{-1}(t) \right)' - \frac{\alpha^2}{\theta^2} B^{-1}(t) \right\}. \quad (2.3)$$

Now we give the main results of this paper.

Theorem 1. Let $\Phi(t, s, r)$ be defined by (1.3). If there exist two positive and real-valued functions $\phi, \theta \in C^1[t_0, \infty)$ such that for every $r \geq t_0$

$$\lim_{t \rightarrow \infty} \sup \lambda_1 \left[\int_r^t M_1(t, s, r) ds \right] > 0, \quad (2.4)$$

where

$$M_1(t, s, r) = \Phi^2(t, s, r)D_1(s) + \Phi'_s(t, s, r)\Phi(t, s, r)K_1(s) - (\Phi'_s(t, s, r))^2 B_1^{-1}(s), \quad (2.5)$$

$$D_1(t) = -C_1(t) - (A^* B_1^{-1} A)(t), \quad K_1(t) = (A^* B_1^{-1} + B_1^{-1} A)(t), \quad (2.6)$$

then (1.1) is oscillatory.

Proof. Assume to the contrary that (1.1) is nonoscillatory. Then $X(t)$ is nonsingular for all sufficiently large t , say $t \geq T \geq t_0$, for any conjoined basis $X(t), Y(t)$ of (1.1). This allows us to make a transformation $W(t) = V(t)U^{-1}(t)$, $t \geq T$. From (2.1) we have

$$\begin{aligned} W'(t) &= -(A^* W + W A + W B_1 W - C_1)(t) \\ &= -\{(W + B_1^{-1} A)^* B_1 (W + B_1^{-1} A)\}(t) - D_1(t). \end{aligned} \quad (2.7)$$

Multiplying (2.7), with t replaced by s , by $\Phi^2(t, s, T)$ and integrating from T to t , we obtain

$$\begin{aligned} \int_T^t \Phi^2(t, s, T) D_1(s) ds &= - \int_T^t \Phi^2(t, s, T) W'(s) ds \\ &\quad - \int_T^t \Phi^2(t, s, T) \{(W + B_1^{-1} A)^* B_1 (W + B_1^{-1} A)\}(s) ds \\ &= 2 \int_T^t \Phi(t, s, T) \Phi'_s(t, s, T) W(s) ds \\ &\quad - \int_T^t \Phi^2(t, s, T) \{(W + B_1^{-1} A)^* B_1 (W + B_1^{-1} A)\}(s) ds. \end{aligned} \quad (2.8)$$

According to the direct computation, we see that

$$\begin{aligned} &2 \int_T^t \Phi(t, s, T) \Phi'_s(t, s, T) W(s) ds \\ &\quad - \int_T^t \Phi^2(t, s, T) \{(W + B_1^{-1} A)^* B_1 (W + B_1^{-1} A)\}(s) ds \end{aligned}$$

$$\begin{aligned}
&= -R_1^{-1}(s)Q_1^*(t, s, T)Q_1(t, s, T)R_1^{-1}(s) \\
&\quad - \Phi(t, s, T)\Phi'_s(t, s, T)K_1(s) + (\Phi'_s(t, s, T))^2B_1^{-1}(s),
\end{aligned}$$

where

$$\begin{aligned}
R_1(s) &= B_1^{1/2}(s), \\
Q_1(t, s, T) &= \Phi(t, s, T)\{R_1(W + B_1^{-1}A)R_1\}(s) - \Phi'_s(t, s, T)I_n.
\end{aligned}$$

Thus from (2.8) and the above computation, we have

$$\begin{aligned}
&\int_T^t \Phi^2(t, s, T)D_1(s)ds \\
&= -\int_T^t R_1^{-1}(s)Q_1^*(t, s, T)Q_1(t, s, T)R_1^{-1}(s)ds \\
&\quad - \int_T^t [\Phi(t, s, T)\Phi'_s(t, s, T)K_1(s) - (\Phi'_s(t, s, T))^2B_1^{-1}(s)]ds. \tag{2.9}
\end{aligned}$$

Therefore, from (2.5), (2.6), and (2.9), for $t \geq T$ we obtain

$$\int_T^t M_1(t, s, T)ds = -\int_T^t R_1^{-1}(s)Q_1^*(t, s, T)Q_1(t, s, T)R_1^{-1}(s)ds \leq 0. \tag{2.10}$$

This implies that

$$\lim_{t \rightarrow \infty} \sup \lambda_1 \left[\int_T^t M_1(t, s, T)ds \right] \leq 0,$$

which contradicts (2.4). This completes the proof of Theorem 1. \square

Remark. We observe that only the ratios α/θ and ϕ/θ are involved in the coefficients of (2.2) and (2.3), and $\alpha(t) = \theta\{(1/2)\log[\phi(t)/\theta(t)]\}'$. Therefore, if we put

$$a(t) = \frac{\theta(t)}{\phi(t)} = \exp\left(-2 \int_{t_0}^t f(s)ds\right),$$

where $f \in C^1[t_0, \infty)$, then

$$\frac{\alpha(t)}{\theta(t)} = f(t), \quad \frac{\alpha(t)}{\phi(t)} = a(t)f(t),$$

and hence

$$\begin{aligned}
B_1(t) &= \frac{1}{a(t)}B(t), \\
C_1(t) &= a(t)[C(t) + f(t)(B^{-1}A + A^*B^{-1})(t) + (f(t)B^{-1}(t))' - f^2(t)B^{-1}(t)].
\end{aligned}$$

In other words, to carry out the transformation, we need only to choose one appropriate smooth function $f(t)$.

If we choose appropriate θ and ϕ in Theorem 1 such that $B_1^{-1}(t) = (\theta/\phi)B^{-1}(t) \leq I_n$ for $t \geq t_0$, i.e., $a(t)B^{-1}(t) \leq I_n$ for $t \geq t_0$, and let $\Phi(t, s, r) = (t-s)(s-r)^\alpha$ for $\alpha > 1/2$, then we have the following theorem from Theorem 1.

Theorem 2. System (1.1) is oscillatory provided that for some $\alpha > 1/2$ and for every $r \geq t_0$

$$\limsup_{t \rightarrow \infty} \frac{1}{t^{2\alpha+1}} \lambda_1 \left[\int_r^t (t-s)^2 (s-r)^{2\alpha} \left(D_1(s) + \frac{\alpha t - (\alpha+1)s + r}{(t-s)(s-r)} K_1(s) \right) ds \right] > \frac{\alpha}{(2\alpha-1)(2\alpha+1)}, \quad (2.11)$$

where $D_1(s)$, $K_1(s)$ are defined as in Theorem 1.

Proof. Assume to the contrary that (1.1) is nonoscillatory. Then $X(t)$ is nonsingular for all sufficiently large t , say $t \geq T \geq t_0$. Similar to the proof of Theorem 1 and because of $B_1^{-1} \leq I_n$, for $t \geq T \geq t_0$ we have

$$\begin{aligned} & \int_T^t \{ (t-s)^2 (s-T)^{2\alpha} D_1(s) + (t-s)(s-T)^{2\alpha-1} [\alpha t - (\alpha+1)s + T] K_1(s) \} ds \\ & \leq \int_T^t [\alpha(t-s)(s-T)^{\alpha-1} - (s-T)^\alpha]^2 B_1^{-1}(s) ds \\ & \leq \int_T^t [\alpha(t-s)(s-T)^{\alpha-1} - (s-T)^\alpha]^2 I_n ds. \end{aligned}$$

This implies that

$$\begin{aligned} & \lambda_1 \left[\int_T^t (t-s)^2 (s-T)^{2\alpha} \left(D_1(s) + \frac{\alpha t - (\alpha+1)s + T}{(t-s)(s-T)} K_1(s) \right) ds \right] \\ & \leq \int_T^t [\alpha(t-s)(s-T)^{\alpha-1} - (s-T)^\alpha]^2 ds. \end{aligned} \quad (2.12)$$

Note that for any $T \geq t_0$

$$\int_T^t [\alpha(t-s)(s-T)^{\alpha-1} - (s-T)^\alpha]^2 ds = \frac{\alpha}{(2\alpha-1)(2\alpha+1)} (t-T)^{2\alpha+1}. \quad (2.13)$$

Thus, from (2.12) and (2.13) we obtain that

$$\begin{aligned} & \lambda_1 \left[\int_T^t (t-s)^2 (s-T)^{2\alpha} \left(D_1(s) + \frac{\alpha t - (\alpha+1)s + T}{(t-s)(s-T)} K_1(s) \right) ds \right] \\ & \leq \frac{\alpha}{(2\alpha-1)(2\alpha+1)} (t-T)^{2\alpha+1}. \end{aligned}$$

It follows that

$$\begin{aligned} & \lim_{t \rightarrow \infty} \sup \frac{1}{t^{2\alpha+1}} \lambda_1 \left[\int_r^t (t-s)^2 (s-T)^{2\alpha} \left(D_1(s) + \frac{\alpha t - (\alpha+1)s + T}{(t-s)(s-T)} K_1(s) \right) ds \right] \\ & \leq \frac{\alpha}{(2\alpha-1)(2\alpha+1)}, \end{aligned}$$

which contradicts assumption (2.11). This completes the proof of Theorem 2. \square

If we choose $\Phi(t, s, r) = (t-s)^\alpha (s-r)$ for $\alpha > 1/2$, similar to the proof of Theorem 2, we can easily obtain the following theorem.

Theorem 3. System (1.1) is oscillatory provided that for some $\alpha > 1/2$ and for every $r \geq t_0$

$$\begin{aligned} & \lim_{t \rightarrow \infty} \sup \frac{1}{t^{2\alpha+1}} \lambda_1 \left[\int_r^t (t-s)^{2\alpha} (s-r)^2 \left(D_1(s) + \frac{t - (\alpha+1)s + \alpha r}{(t-s)(s-r)} K_1(s) \right) ds \right] \\ & > \frac{\alpha}{(2\alpha-1)(2\alpha+1)}, \end{aligned} \quad (2.14)$$

where $D_1(s)$, $K_1(s)$ are defined as in Theorem 1.

Proof. Assume to the contrary that (1.1) is nonoscillatory. Then $X(t)$ is nonsingular for all sufficiently large t , say $t \geq T \geq t_0$. Similar to the proof of Theorem 2 and because of $B_1^{-1} \leq I_n$, for $t \geq T \geq t_0$ we have

$$\begin{aligned} & \lambda_1 \left[\int_T^t (t-s)^{2\alpha} (s-T)^2 \left(D_1(s) + \frac{t - (\alpha+1)s + \alpha T}{(t-s)(s-T)} K_1(s) \right) ds \right] \\ & \leq \int_T^t [(t-s)^\alpha - \alpha(t-s)^{\alpha-1}(s-T)]^2 ds. \end{aligned}$$

Noting that

$$\int_T^t [(t-s)^\alpha - \alpha(t-s)^{\alpha-1}(s-T)]^2 ds = \frac{\alpha}{(2\alpha-1)(2\alpha+1)} (t-T)^{2\alpha+1},$$

we obtain a contradiction that

$$\lim_{t \rightarrow \infty} \sup \frac{1}{t^{2\alpha+1}} \lambda_1 \left[\int_T^t (t-s)^{2\alpha} (s-T)^2 \left(D_1(s) + \frac{t - (\alpha+1)s + \alpha T}{(t-s)(s-T)} K_1(s) \right) ds \right] \leq \frac{\alpha}{(2\alpha-1)(2\alpha+1)}.$$

This completes the proof of Theorem 3. \square

In order to show the sharpness of our results, now let us consider the following two examples:

Example 1. Consider the Euler differential system

$$Y'' + \text{diag}\left(\frac{\beta}{t^2}, \frac{\gamma}{t^2}\right)Y = 0, \quad t \geq 1, \quad \gamma \geq \beta > 0. \quad (2.15)$$

If we choose $f(t) = 0$, then $a(t) = 1$, $D_1(t) = \text{diag}(\beta/t^2, \gamma/t^2)$ and $K_1(t) = 0$. Note that for each $r \geq t_0$

$$\lim_{t \rightarrow \infty} \frac{1}{t^{2\alpha+1}} \int_r^t (t-s)^2 (s-r)^{2\alpha} \frac{\gamma}{s^2} ds = \frac{\gamma}{\alpha(2\alpha-1)(2\alpha+1)}.$$

For any $\gamma > 1/4$, there exists $\alpha > 1/2$ such that

$$\frac{\gamma}{\alpha(2\alpha-1)(2\alpha+1)} > \frac{\alpha}{(2\alpha-1)(2\alpha+1)},$$

i.e., $\gamma > \alpha^2$. This means that (2.11) holds. By Theorem 2, we find that system (2.15) is oscillatory for $\gamma > 1/4$. However, we can easily show that criteria in [3,6,8] fail to reveal this fact.

Example 2. Consider the 4-dimensional system (1.1) with $t \geq 1$ and

$$A(t) = \begin{bmatrix} 0 & -1/t \\ 2/t & 0 \end{bmatrix}, \quad B(t) = \begin{bmatrix} t & 0 \\ 0 & 2t \end{bmatrix}, \quad C(t) = -\begin{bmatrix} \theta/t^3 & 0 \\ 0 & \eta/t^3 \end{bmatrix},$$

where $\eta \geq \theta > 0$ and $t \geq 1$. According to the remark, if we let $f(t) = -1/2t$, then $a(t) = t$ and $a(t)B^{-1}(t) \leq I_2$ for $t \geq 1$. Thus, from (2.6) we have

$$K_1(t) \equiv 0, \quad D_1(t) = \begin{bmatrix} (\theta - 11/4)t^{-2} & 0 \\ 0 & (\eta - 11/8)t^{-2} \end{bmatrix}.$$

Similar to the proof of Example 1, we can obtain that system (1.1) is oscillatory when $\eta > 13/8$ by Theorem 2.

References

- [1] G.J. Butler, L.H. Erbe, Oscillation results for second order differential systems, SIAM J. Math. Anal. 17 (1986) 19–29.

- [2] G.J. Butler, L.H. Erbe, Oscillation results for self-adjoint differential systems, *J. Math. Anal. Appl.* 115 (1986) 470–481.
- [3] G.J. Butler, L.H. Erbe, A.B. Mingarelli, Riccati techniques and variational principles in oscillation theory for linear systems, *Trans. Amer. Math. Soc.* 303 (1987) 263–282.
- [4] R. Byers, B.J. Harris, M.K. Kwong, Weighted means and oscillation conditions for second order matrix differential equations, *J. Differential Equations* 61 (1986) 164–177.
- [5] W.A. Coppel, Disconjugacy, in: *Lecture Notes in Mathematics*, Vol. 220, Springer-Verlag, Berlin, 1971.
- [6] L.H. Erbe, Q. Kong, S. Ruan, Kamenev type theorems for second order matrix differential systems, *Proc. Amer. Math. Soc.* 117 (1993) 957–962.
- [7] P. Hartman, Self-adjoint, non-oscillatory systems of ordinary second order, linear differential equations, *Duke Math. J.* 24 (1957) 25–36.
- [8] D.B. Hinton, R.T. Lewis, Oscillation theory for generalized second order differential equations, *Rocky Mountain J. Math.* 10 (1980) 751–766.
- [9] I. Sowjaya Kumari, S. Umanaheswaram, Oscillation criteria for linear matrix Hamiltonian systems, *J. Differential Equations* 165 (2000) 174–198.
- [10] M.K. Kwong, H.G. Kaper, Oscillation of two-dimensional linear second order differential systems, *J. Differential Equations* 56 (1985) 195–205.
- [11] M.K. Kwong, H.G. Kaper, K. Akiyama, A.B. Mingarelli, Oscillation of linear second-order differential systems, *Proc. Amer. Math. Soc.* 91 (1984) 85–91.
- [12] F. Meng, J. Wang, Z. Zheng, A note on Kamenev type theorems for second order matrix differential systems, *Proc. Amer. Math. Soc.* 126 (1998) 391–395.
- [13] A.B. Mingarelli, On a conjecture for oscillation of second order ordinary differential systems, *Proc. Amer. Math. Soc.* 82 (1981) 593–598.
- [14] A. Peterson, J. Ridenhour, Oscillation of second order linear matrix difference equations, *J. Differential Equations* 89 (1991) 69–88.